

CONIFEROUS-HABITAT ASSOCIATIONS OF SPIDERS (ARANEAE) ON RED SPRUCE FOLIAGE

Daniel T. Jennings

Northeastern Forest Experiment Station
USDA Building, University of Maine
Orono, Maine 04469

and

Judith A. Collins

Department of Entomology
University of Maine
Orono, Maine 04469

ABSTRACT

Coniferous-habitat associations were determined for 16 species of adult spiders collected from red spruce foliage in northern Maine. Most of the spiders have been found on three or more (range one to 15) conifer species. Spider-conifer habitats were positively correlated ($r = 0.96$) with geographic states and provinces. Mean conifer-habitat associations did not differ between web spinner and hunter species. Significantly more species of spiders from red spruce foliage were associated with northern conifers than with western conifers, but not with southern conifers.

Comparison of the red spruce fauna with 16 selected spider-tree faunal studies showed that 1) Sørensen's similarity quotients (QS) were significantly greater for genera than for species, and 2) mean species QS values were significantly greater for northern than for southern but not for western conifers. The spider fauna on red spruce is closely allied with that on other northern conifers; however, none of the 16 spider species are restricted to conifers.

INTRODUCTION

The arboreal spider fauna of North American trees has received little attention; fewer than 25 coniferous and deciduous tree species have been studied in detail (Jennings 1976). Recent studies include: spiders on red pine, *Pinus resinosa* Ait., white spruce, *Picea glauca* (Moench) Voss, and northern white-cedar, *Thuja occidentalis* L., in Minnesota (Stratton, Uetz, and Dillery 1979); spiders on white fir, *Abies concolor* (Gord. and Glend.) Lindl. ex Hildebr., in California (Dahlsten et al. 1977) and in Oregon (Mason and Torgersen 1983). Additional spider-tree association records are scattered throughout the araneological and entomological literature; most records are incidental to other study objectives.

During our investigations on natural enemies of the spruce budworm, *Choristoneura fumiferana* (Clem.), we collected spiders of 10 families, 16 genera, and 21 species from foliage of red spruce, *Picea rubens* Sarg., in northern Maine. Mean spider densities during two sampling periods and spider-budworm

relationships were discussed in an earlier paper (Jennings and Collins 1987). Here we summarize the known coniferous-habitat associations for each of the 16 species found on red spruce foliage, and compare the spider fauna on red spruce with that of other coniferous species in North America. Our purposes for these comparisons were: 1) to identify (by spider and conifer species) the known spider-coniferous habitat associations, 2) to determine possible relationships between spider-coniferous habitats and geographic states and provinces, 3) to determine the degree of similarity between the spider fauna on red spruce and that found on other conifers, and 4) to determine possible patterns of association or differences in association among geographic regions. This information may be useful to forest pest managers who are concerned with identifying potentially important predators of coniferous pests.

METHODS

Habitat Associations.—We searched the araneological and entomological literature for spider-tree faunal studies and for habitat-association records of spiders collected from North American conifers. Standard reference sources were consulted including Biological Abstracts, Zoological Record, and Centre International de Documentation Arachnologique Liste. In addition, a DIALOG search was made from the AGRICOLA database at the National Agricultural Library, Beltsville, Maryland. The database search was limited to North American literature that included information and identities of both spiders and trees.

Because tree habitats may vary by spider species, and spiders identified to generic level only may include numerous species, we restricted our search to the 16 species of adult spiders found on red spruce foliage (Jennings and Collins 1987). In searching the older literature, recent spider synonymies were considered, e.g., *Pityohyphantes phrygianus* (Koch) [= *P. costatus* (Hentz)] (Kaston 1981), *Neoscona minima* F.O.P.-Cambridge [= *N. arabesca* (Walck.)] (Berman and Levi 1971), and *Araneus displicata* (Hentz) [= *Araniella displicata* (Hentz)] (Levi 1974). Consistent with Dondale (1959), records of *Grammonota pictilis* (O. P. Cambridge) collected on balsam fir, *Abies balsamea* (L.) Mill., in New Brunswick were considered to belong to *G. angusta* Dondale.

Habitat associations were recorded by spider and tree species. Only coniferous tree species were considered, though many of the red spruce spiders also occur on broad-leaved deciduous trees. Common and scientific names of trees follow Little (1979). Both evergreen and cone-bearing species of the families Cupressaceae, Pinaceae, and Taxodiaceae were included.

Data Analyses.—Although our literature search did not represent a random sample (i.e., all known available studies were included), we considered the data generated by these faunal studies to meet random-sample criteria (i.e., all potential coniferous habitats were available for study; none were selectively biased for our comparison purposes). We used nonparametric procedures (Sokal and Rohlf 1981) for statistical comparisons. The Wilcoxon two-sample test was used to test for differences between and among means ($P = 0.05$). For comparisons involving more than two means, all 2-mean combinations were performed.

Correlation analysis (Sokal and Rohlf 1981) was used as a measure of association between spider-coniferous habitats and geographic states and

provinces. We defined a spider-coniferous habitat as the collection of a spider species from or on a conifer species, e.g., *Dictyna brevitarus* Emerton collected from foliage of *Abies balsamea* (L.) Mill. in New Brunswick (Renault 1968). Regression analysis was not used because we were interested in the degree of association (interdependence) between the two variables, not the dependence of one on the other. Habitat records without geographic locality were excluded from the analysis.

Sørensen's similarity quotient (QS), as defined by Price (1975), was used to determine the degree of similarity between the spider fauna on red spruce and that found on other coniferous species. The formula used was: $QS = 2c \times 100 / (a + b)$, where a = the number of spider genera or species in study A; b = the number of spider genera or species in study B; and c = the number of spider genera or species common to both studies.

To determine possible patterns of association among geographic areas, each conifer species was assigned to one of three broadly defined regions—northern, southern, or western—based on distributional ranges of trees given by Little (1979). We then calculated mean QS values for each region and performed statistical tests to determine possible differences and associations among regions. Likewise, mean numbers of spider species in common with red spruce were determined and compared for each region.

RESULTS AND DISCUSSION

Coniferous-Habitat Associations.—The 16 species of adult spiders found on red spruce foliage and their habitat associations with North American conifers are summarized in Table 1. With only one exception, *Xysticus discursans* Keyserling, most of the spiders have been collected from three or more conifer species (range 1 to 15). Included are habitat affinities with balsam fir, *Abies balsamea* (L.) Mill., and white fir, *A. concolor* (Gord. and Glend.) Lindl. ex Hildebr.; oldfield common juniper, *Juniperus communis* var. *depressa* Pursh., and *Juniperus* spp. (may include *J. chinensis* L., eastern red cedar, *J. virginiana* L., and Rocky Mountain juniper, *J. scopulorum* Sarg.); tamarack, *Larix laricina* (Du Roi) K. Koch, and western larch, *L. occidentalis* Nutt.; Norway spruce, *Picea abies* (L.) Karst., white spruce, *P. glauca* (Moench) Voss, and red spruce, *P. rubens* Sarg.; jack pine, *Pinus banksiana* Lamb., sand pine, *P. clausa* (Chapm. ex Engelm.) Vasey ex Sarg., lodgepole pine, *P. contorta* var. *latifolia* Engelm., shortleaf pine, *P. echinata* Mill., Jeffrey pine, *P. jeffreyi* Grev. and Balf., ponderosa pine, *P. ponderosa* Dougl. ex Laws., red pine, *P. resinosa* Ait., eastern white pine, *P. strobus* L., Scotch pine, *P. sylvestris* L., loblolly pine, *P. taeda* L., and Virginia pine, *P. virginiana* Mill.; Douglas-fir, *Pseudotsuga menziesii* (Mirb.) Franco; baldcypress, *Taxodium* sp.; and northern white-cedar, *Thuja occidentalis* L.

Predictably, coniferous-habitat associations were positively and highly correlated ($r = 0.96$) with geographic states and provinces (Fig. 1). This indicates that, within limits, the more localities sampled, the greater the likelihood of finding more coniferous-habitat associations. However, numbers of available coniferous species vary among states and provinces, and some spider species have limited ranges.

Considering habitat association by spider foraging group indicates that web spinners have been found and reported from more species of conifers ($\bar{x} = 7.9$

Table 1.—Coniferous-habitat associations of spiders from red spruce foliage.

Conifer species	Locality	Reference(s)
<i>Dictyna brevitarus</i> Emerton		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963), Renault (1968) Renault and Miller (1972)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
<i>Pinus strobus</i>	Wisconsin	Coppel and Smythe (1963)
<i>Theridion montanum</i> Emerton		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963), Renault (1968) Renault and Miller (1972)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
	New Mexico	Levi (1957)
<i>Pinus ponderosa</i>	New Mexico	Levi (1957)
<i>Theridion murarium</i> Emerton		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963), Renault (1968) Renault and Miller (1972)
<i>Abies concolor</i>	Oregon	Mason and Torgersen (1983)
<i>Juniperus communis</i> var. <i>depressa</i>	Michigan	Drew (1967)
<i>Juniperus</i> spp.	Kansas	Heinrichs and Thompson (1968)
	New Mexico	Levi (1957)
<i>Picea abies</i>	Massachusetts	Taylor (1928) ¹
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
	Connecticut	Kaston (1981)
<i>Pinus banksiana</i>	Michigan	Allen et al. (1970)
	Indiana	Lowrie (1948)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971) ¹
<i>Pinus ponderosa</i>	New Mexico	Levi (1957)
<i>Pinus resinosa</i>	Ontario	Martin (1966)
<i>Pinus strobus</i>	Wisconsin	Coppel and Smythe (1963)
	Massachusetts	Taylor (1928)
<i>Pinus sylvestris</i>	Massachusetts	Taylor (1928)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971)
	Oklahoma	Bosworth et al. (1971)
<i>Pinus virginiana</i>	Maryland	Howden and Vogt (1951)
<i>Pinus</i> sp.	Connecticut	Kaston (1981)
<i>Pseudotsuga menziesii</i>	British Columbia	Turnbull (1956)
<i>Thuja occidentalis</i>	Michigan	Drew (1967)
<i>Pityohyphantes costatus</i> (Hentz)		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963), Renault (1968) Renault and Miller (1972)
<i>Abies concolor</i>	Oregon	Mason and Torgersen (1983)
	California	Ohmart and Dahlsten (1979)
<i>Picea glauca</i>	Quebec	Manuel (1984)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
<i>Pinus contorta</i> var. <i>latifolia</i>	Alberta	Powell (1971)
<i>Pinus resinosa</i>	Minnesota	Stratton et al. (1979)
	Connecticut	Bean and Godwin (1955)
<i>Thuja occidentalis</i>	Michigan	Drew (1967)

Table 1.—Continued.

<i>Grammonota angusta</i> Dondale		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963), Renault (1968) Renault and Miller (1972)
<i>Larix laricina</i>	Manitoba	Ives (1967)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
<i>Pinus banksiana</i>	Manitoba	Bradley and Hinks (1968)
<i>Eustala anastera</i> (Walckenaer)		
<i>Abies balsamea</i>	New Brunswick	Levi (1977)
<i>Juniperus</i> spp.	Kansas	Heinrichs and Thompson (1968)
	Nebraska	Worley and Pickwell (1927)
<i>Larix occidentalis</i>		Levi (1977)
<i>Picea glauca</i>	New Brunswick	Levi (1977)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Pinus banksiana</i>	Michigan	Allen et al. (1970)
<i>Pinus clausa</i>	Florida	Levi (1977)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971), Levi (1977)
<i>Taxodium</i> sp.		Levi (1977)
<i>Neoscona arabesca</i> (Walckenaer)		
<i>Abies balsamea</i>	New Brunswick	Renault (1968)
<i>Abies concolor</i>	Oregon	Mason and Torgersen (1983)
<i>Juniperus communis</i> var. <i>depressa</i>	Michigan	Drew (1967)
<i>Juniperus</i> sp.		Berman and Levi (1971)
<i>Larix laricina</i>		Berman and Levi (1971)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971)
<i>Pinus strobus</i>	Wisconsin	Coppel and Smythe (1963)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971)
	Oklahoma	Bosworth et al. (1971)
<i>Pinus virginiana</i>	Maryland	Howden and Vogt (1951)
<i>Araniella displicata</i> (Hentz)		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963) Renault and Miller (1972)
<i>Juniperus communis</i> var. <i>depressa</i>	Michigan	Drew (1967)
<i>Picea glauca</i>	Minnesota	Stratton et al. (1979)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
<i>Pinus banksiana</i>	Manitoba	Bradley and Hinks (1968)
	Michigan	Allen et al. (1970)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971)
<i>Pinus jeffreyi</i>	California	Dahlsten (1961) ¹
<i>Pinus ponderosa</i>	California	Dahlsten (1961)
<i>Pinus resinosa</i>	Ontario	Martin (1966)
<i>Pinus strobus</i>	Wisconsin	Coppel and Smythe (1963)
	Maine	Procter (1946)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971)
	Oklahoma	Bosworth et al. (1971)
<i>Pseudotsuga menziesii</i>	British Columbia	Turnbull (1956)
<i>Clubiona canadensis</i> Emerton		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963) Renault (1968) Renault and Miller (1972)

Table 1.—Continued.

<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
<i>Thuja occidentalis</i>	Michigan	Drew (1967)
<i>Philodromus exilis</i> Banks		
<i>Abies balsamea</i>	New Brunswick	Renault (1968)
<i>Abies</i> sp.		Dondale and Redner (1968)
<i>Juniperus</i> sp.		Dondale and Redner (1968)
<i>Picea glauca</i>	Ontario	Dondale (pers. comm.)
	Nova Scotia	
	New Brunswick	
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Pinus strobus</i>	Ontario	Dondale (pers. comm.)
	Nova Scotia	
<i>Thuja occidentalis</i>	Ontario	Dondale (pers. comm.)
<i>Philodromus placidus</i> Banks		
<i>Abies balsamea</i>	New Brunswick	Loughton et al. (1963)
		Renault (1968)
		Renault and Miller (1972)
<i>Juniperus communis</i> var. <i>depressa</i>	Michigan	Drew (1967)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
<i>Pinus banksiana</i>	Manitoba	Bradley and Hinks (1968)
	Michigan	Allen et al. (1970)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971)
<i>Pinus resinosa</i>	Minnesota	Heimer et al. (1984)
<i>Pinus</i> sp.	Maine	Procter (1946)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971)
<i>Misumena vatia</i> (Clerck)		
<i>Juniperus communis</i> var. <i>depressa</i>	Michigan	Drew (1967)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Pinus banksiana</i>	Manitoba	Bradley and Hinks (1968)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971)
<i>Pseudotsuga menziesii</i>	British Columbia	Turnbull (1956)
<i>Xysticus discursans</i> Keyserling		
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Xysticus punctatus</i> Keyserling		
<i>Abies balsamea</i>	New Brunswick	Renault (1968)
<i>Picea glauca</i>	Minnesota	Houseweart and Kulman (1976)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Pinus banksiana</i>	Manitoba	Bradley and Hinks (1968)
	Michigan	Allen et al. (1970)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971)
<i>Pinus resinosa</i>	Minnesota	Heimer et al. (1984)
<i>Pinus strobus</i>	Wisconsin	Coppel and Smythe (1963)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971)
	Oklahoma	Bosworth et al. (1971)
<i>Pseudotsuga menziesii</i>	British Columbia	Turnbull (1956)
<i>Salticus scenicus</i> (Linnaeus)		
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Pinus echinata</i>	North Carolina	Ramsey (1941) ¹
<i>Pinus taeda</i>	North Carolina	Ramsey (1941)
	Oklahoma	Bosworth et al. (1971)

Table 1.—Continued.

	<i>Metaphidippus flavipedes</i> (G. & E. Peckham)	
<i>Abies balsamea</i>	Maine	Jennings and Houseweart (1978)
	New Brunswick	Loughton et al. (1963)
		Renault (1968)
<i>Juniperus communis</i> var. <i>depressa</i>	Michigan	Renault and Miller (1972)
		Drew (1967)
<i>Picea glauca</i>	Minnesota	Houseweart and Kulman (1976)
<i>Picea rubens</i>	Maine	Jennings and Collins (1987)
<i>Picea</i> sp.	New Brunswick	Renault and Miller (1972)
<i>Pinus banksiana</i>	Manitoba	Bradley and Hinks (1968)
	Michigan	Allen et al. (1970)
<i>Pinus echinata</i>	Arkansas	Peck et al. (1971)
<i>Pinus resinosa</i>	Ontario	Martin (1966)
	Minnesota	Heimer et al. (1984)
<i>Pinus strobus</i>	Wisconsin	Coppel and Smythe (1963)
<i>Pinus taeda</i>	Arkansas	Peck et al. (1971)
<i>Pseudotsuga menziesii</i>	British Columbia	Turnbull (1956)
<i>Thuja occidentalis</i>	Michigan	Drew (1967)

¹Collections not separated by tree species.


± 1.6) than have hunters ($\bar{x} = 5.8 \pm 1.2$); however, the means for these two groups were not significantly different by the Wilcoxon two-sample test. Coniferous foliage is a suitable habitat substrate for both spider groups. Stratton et al. (1979) found greater percentages of web-builders than hunters on pine, spruce, and cedar in Minnesota; but the relative composition of the two foraging groups was not significantly different among tree species.

Exotic conifers excepted, i.e., *Picea abies* and *Pinus sylvestris*, significantly more ($P = 0.02$) species of spiders from red spruce foliage were associated with northern conifers ($\bar{x} = 7.4 \pm 1.5$) than with western conifers ($\bar{x} = 2.9 \pm 0.8$), but not with southern conifers ($\bar{x} = 4.4 \pm 1.9$). This supports our hypothesis that the spider fauna on red spruce is closely allied with other northern conifers; however, elements may also occur on southern conifers, and to a lesser degree on western conifers. Mean numbers of red spruce species associated with western and southern conifers did not differ significantly.

Among individual species, *Theridion murarium* Emerton showed the greatest range of coniferous habitats, being found on at least 15 species of conifers. Six other spider species have been collected from seven or more conifer species: *Araniella displicata* (Hentz) (12), *Metaphidippus flavipedes* (G. & E. Peckham) (11), *Xysticus punctatus* Keyserling (9), *Eustala anastera* (Walckenaer) (8), *Neoscona arabesca* (Walckenaer) (8), and *Pityohyphantes costatus* (Hentz) (7). The remaining spiders from red spruce foliage are reported from six or fewer conifers. Apparently *Xysticus discursans* has been collected and reported from only one conifer, *Picea rubens*; however, this association may be accidental because the normal habitat for this spider is near the ground (Dondale and Redner 1978).

The number of coniferous-habitat associations for individual spider species does not necessarily correspond with or indicate their habitat specificity. None of the 16 species found on red spruce are restricted to that tree species, or to

Fig. 1.—Relationship between spider-coniferous habitats and states and provinces; correlation analysis, $r = 0.96$ (Sokal and Rohlf 1981). Each point represents the number of conifer species and states and provinces from which one of the red spruce spiders ($n = 16$) has been reported.



conifers. For example, many of the species, including *Theridion murarium*, *Metaphidippus flavipedes*, *Araniella displicata*, *Eustala anastera*, and *Neoscona arabesca*, also occur on a variety of broad-leaved trees and shrubs. *Misumena vatia* (Clerck) and *Clubiona canadensis* Emerton are frequently collected from shrubs and forbs; *Salticus scenicus* (Linnaeus) is found in synanthropic habitats (Kaston 1981). *Xysticus discursans* is usually taken by pitfall traps and sweep nets in both grassland and wooded areas (Dondale and Redner 1978). *Grammonota angusta* Dondale, *Philodromus placidus* Banks, and *Xysticus punctatus* are typically found on conifers.

Spider-Faunal Studies.—Sørensen's similarity quotients (QS values) for 16 spider-tree faunal studies are shown in Table 2. Not included are studies lacking complete species lists (Dahlsten et al. 1977; Renault and Miller 1972; Stratton et al. 1979) and studies where tree-habitat association was uncertain (Fox and Griffith 1976). As expected, QS values were significantly greater ($P < 0.01$) for genera than for species, i.e., more genera than species were shared in common among the studies. QS values for species were generally < 30 , no doubt because sampling methods and intensities varied considerably among the studies. However, despite these differences, the red spruce fauna showed more similarities with spider faunas on northern conifers than on southern or western conifers. Mean species QS values were significantly greater ($P = 0.03$) for northern conifers ($\bar{x} = 18.4 \pm 2.2$) than for southern conifers ($\bar{x} = 9.0 \pm 1.5$), but not for western conifers ($\bar{x} = 13.2 \pm 0.8$). Southern and western QS means also did not differ significantly; however, the level of statistical significance ($P = 0.08$) shows possible distinction. With only two exceptions, *Larix laricina* and *Pinus resinosa*, QS values for northern conifers were generally > 15 ; whereas, those for southern and western conifers were generally < 15 .

The apparent similarity among spider faunas on northern conifers is also evidenced by comparable QS values, particularly for tree species often found growing in the same forest stand, e.g., *Abies balsamea*, *Pinus strobus*, and *Thuja occidentalis* (Table 2). Surprisingly, *Larix laricina*, a common resident of northeastern spruce-fir forests, had a very low QS value; conversely, *Juniperus communis* var. *depressa*, an inhabitant of old fields and cutover forests, had a relatively high QS value. We are unable to explain these dissimilarities.

Table 2.—Comparison of red spruce spider fauna with spider faunas of other coniferous species by geographic region. QS = Sørensen's similarity quotient.

Conifer species	State- Province	QS value		Reference
		Genera	Species	
NORTHERN REGION				
<i>Abies balsamea</i>	New Brunswick	30.4	19.5	Renault (1968)
<i>Abies balsamea</i>	New Brunswick	37.7	24.0	Loughton et al. (1963)
<i>Juniperus communis</i> var. <i>depressa</i>	Michigan	57.1	27.8	Drew (1967)
<i>Larix laricina</i>	Manitoba	51.8	6.1	Ives (1967)
<i>Pinus banksiana</i>	Michigan	46.2	21.4	Bradley and Hinks (1968)
<i>Pinus banksiana</i>	Michigan	38.5	18.8	Allen et al. (1970)
<i>Pinus resinosa</i>	Ontario	25.8	10.5	Martin (1966)
<i>Pinus strobus</i>	Wisconsin	59.5	19.4	Coppel and Smythe (1963)
<i>Thuja occidentalis</i>	Michigan	38.5	18.2	Drew (1967)
SOUTHERN REGION				
<i>Juniperus</i> spp.	Kansas	37.5	9.5	Heinrichs and Thompson (1968)
<i>Pinus echinata</i> and <i>P. taeda</i> ¹	North Carolina	25.9	3.2	Ramsey (1941)
<i>Pinus echinata</i> and <i>P. taeda</i> ¹	Arkansas	27.7	10.3	Peck et al. (1971)
<i>Pinus taeda</i>	Oklahoma	32.4	11.6	Bosworth et al. (1971)
<i>Pinus virginiana</i>	Maryland	32.3	10.5	Howden and Vogt (1951)
WESTERN REGION				
<i>Abies concolor</i>	Oregon	54.6	14.0	Mason and Torgersen (1983)
<i>Pseudotsuga menziesii</i>	British Columbia	36.7	12.5	Turnbull (1956)

¹Collections not separated by tree species.

In addition to geographic region, tree height and growth form may also influence resident spider faunas. For example, within the same northern region, spider faunas on young, plantation red pine and on tamarack generally had low QS values compared with other species. Martin (1966) concluded that young red pine seedlings planted in an old field do not form an influential part of the community, and ecological conditions remain those of an old field. No doubt the sparse, clumped needles of tamarack provide fewer microhabitats for spiders than the dense, dispersed foliage of spruce. Stratton et al. (1979) found the highest number of spider species and individuals on foliage of white spruce compared with red pine and northern white-cedar.

Notably absent from these faunistic comparisons, including our own study, are investigations involving two or more coniferous species sampled at the same time, place, and intensity. For such studies, we predict that QS values will be much higher, particularly for conifers growing in the same forest stand and of similar growth form, height, and age. Under these conditions, faunal similarities may approach the QS-50 limit, below which communities are arbitrarily considered distinct (Price 1975). None of the species faunistic comparisons in Table 2 approached this limit, probably because the conifers studied were widely separated in time, place, and sampling methodology.

Finally, the paucity of information about spider-coniferous habitat associations is evident from both Tables 1 and 2. Apparently less than 30% of the 82 native species of conifers (Little 1979; families Cupressaceae, Pinaceae, and Taxodiaceae) have even been examined for spiders, much less studied in detail. The maximum

number of conifer species sampled in any one state or province was four; likewise, only one conifer, *Picea glauca*, has been studied in five states and provinces. We conclude that additional faunistic studies are needed: 1) to define the spider fauna of any one conifer species, and 2) to determine habitat specificities and associational ranges of individual spider species. Such information will help to elucidate potential predator-prey relationships involving spiders and pests of conifers.

ACKNOWLEDGMENTS

We thank Daniel Starr, who at the time of this study was with USDA Forest Service, National Agricultural Library, Beltsville, Maryland, for assistance with literature searches. Richard A. Hosmer provided computer programming assistance and is with the USDA Forest Service, Northeastern Forest Experiment Station, Orono, Maine.

We are grateful to our reviewers, Bruce Cutler, Charles D. Dondale, James H. Redner, and Dave Walter for their constructive comments, and to Donald W. Seegrist, Biometrician, Northeastern Forest Experiment Station, for statistical review.

Portions of this research were supported by funds provided by the USDA, Forest Service, Northeastern Forest Experiment Station, NE-1151, to the Department of Entomology, College of Life Sciences and Agriculture, University of Maine, Orono (Cooperative Research Agreement 23-805).

LITERATURE CITED

- Allen, D. C., F. B. Knight, and J. L. Foltz. 1970. Invertebrate predators of the jack-pine budworm, *Choristoneura pinus*, in Michigan. Ann. Entomol. Soc. Amer., 63:59-64.
- Bean, J. L. and P. A. Godwin. 1955. Description and bionomics of a new red pine scale, *Matsucoccus resinosa*. For. Sci., 1:164-176.
- Berman, J. D. and H. W. Levi. 1971. The orb weaver genus *Neoscona* in North America. Mus. Comp. Zool. Bull., 5:465-500.
- Bosworth, A. B., H. G. Raney, E. D. Sturgeon, R. D. Morrison, and R. D. Eikenbary. 1971. Population trends and location of spiders in loblolly pines, with notes of predation on the *Rhyacionia* complex (Lepidoptera: Olethreutidae). Ann. Entomol. Soc. Amer., 64:864-870.
- Bradley, G. A. and J. D. Hinks. 1968. Ants, aphids, and jack pine in Manitoba. Canadian Entomol., 100:40-50.
- Coppel, H. C. and R. V. Smythe. 1963. Occurrence of spiders on eastern white pine trees infested with the introduced pine sawfly, *Diprion similis* (Htg.). Univ. Wisconsin Forestry Res. Notes No. 102. 7 pp.
- Dahlsten, D. L. 1961. Life history of a pine sawfly, *Neodiprion* sp., at Willits, California (Hymenoptera: Diprionidae). Canadian Entomol., 93:182-195.
- Dahlsten, D. L., R. F. Luck, E. I. Schlinger, J. M. Wenz., and W. A. Copper. 1977. Parasitoids and predators of the Douglas-fir tussock moth, *Orgyia pseudotsugata* (Lepidoptera: Lymantriidae), in low to moderate populations in central California. Canadian Entomol., 109:727-746.
- Dondale, C. D. 1959. Definition of the genus *Grammonota* with descriptions of seven new species. Canadian Entomol., 91:232-242.
- Dondale, C. D. and J. H. Redner. 1968. The *imbecillus* and *rufus* groups of the spider genus *Philodromus*. Entomol. Soc. Canada Mem., 55. 78 pp.
- Dondale, C. D. and J. H. Redner. 1978. The crab spiders of Canada and Alaska (Araneae: Philodromidae and Thomisidae). Canadian Dept. Agr. Publ., 1663. 255 pp.

- Drew, L. C. 1967. Spiders of Beaver Island Michigan. Publications of the Museum, Michigan State Univ., 3(3):153-208.
- Fox, R. C. and K. H. Griffith. 1976. Predation of pine cinaran aphids by spiders. J. Georgia Entomol. Soc., 11:241-243.
- Heinrichs, E. A. and H. E. Thompson. 1968. Parasites, predators, and other arthropods associated with *Choristoneura houstonana* (Lepidoptera: Tortricidae) on *Juniperus* species in Kansas. Canadian Entomol., 100:33-39.
- Heimer, S., W. Nentwig, and B. Cutler. 1984. The spider fauna of the Itasca State Park (Minnesota, USA). Faun. Abh. Mus. Tierk. Dresden, II, Nr. 6, 1983:119-124.
- Houseweart, M. W. and H. W. Kulman. 1976. Life tables of the yellowheaded spruce sawfly, *Pikonema alaskensis* (Rohwer) (Hymenoptera: Tenthredinidae) in Minnesota. Environ. Entomol., 5:859-867.
- Howden, H. F. and G. B. Vogt. 1951. Insect communities of standing dead pine (*Pinus virginiana* Mill.). Ann. Entomol. Soc. Amer., 44:581-595.
- Ives, W. G. H. 1967. Relations between invertebrate predators and prey associated with larch sawfly eggs and larvae on tamarack. Canadian Entomol., 99:607-622.
- Jennings, D. T. 1976. Spiders on black walnut. Amer. Midland Nat., 95:111-119.
- Jennings, D. T. and J. A. Collins. 1987. Spiders on red spruce foliage in northern Maine. J. Arachnol., 14:303-313.
- Jennings, D. T. and M. W. Houseweart. 1978. Spider preys on spruce budworm egg mass. Entomol. News, 89:183-186.
- Kaston, B. J. 1981. Spiders of Connecticut. Bull. Connecticut State Geol. Nat. Hist. Surv., 70: 1020 pp.
- Levi, H. W. 1957. The spider genera *Enoplognatha*, *Theridion*, and *Paidisca* in America north of Mexico. Amer. Mus. Nat. Hist. Bull., 112:5-123.
- Levi, H. W. 1974. The orbweaver genera *Araniella* and *Nuctenea*. Mus. Comp. Zool. Bull., 146:291-316.
- Levi, H. W. 1977. The American orbweaver genera *Cyclosa*, *Metazygia* and *Eustala* north of Mexico. Mus. Comp. Zool. Bull., 148:61-127.
- Little, E. L., Jr. 1979. Checklist of United States trees (native and naturalized). For. Serv., U.S. Dep. Agr., Washington, D.C. Agr. Handbk., No. 541. 375 pp.
- Loughton, B. G., C. Derry, and A. S. West. 1963. Spiders and the spruce budworm. Pp. 249-268. In The dynamics of epidemic spruce budworm populations. (R. F. Morris, ed.). Mem. Entomol. Soc. Canada 31. 332 pp.
- Lowrie, D. C. 1948. The ecological succession of the spiders of the Chicago area dunes. Ecology, 29:334-351.
- Manuel, R. L. 1984. The egg sac of *Pityohyphantes costatus* (Hentz) (Araneae, Linyphiidae) and its phorid parasite. J. Arachnol., 12:371-372.
- Mason, R. R. and T. R. Torgersen. 1983. Mortality of larvae in stocked cohorts of the Douglas-fir tussock moth, *Orgyia pseudotsugata* (Lepidoptera: Lymantriidae). Canadian Entomol., 115:1119-1127.
- Martin, J. L. 1966. The insect ecology of red pine plantations in central Ontario. IV. The crown fauna. Canadian Entomol., 98:10-27.
- Ohmart, C. P. and D. L. Dahlsten. 1979. Biological studies of bud mining sawflies, *Pleroneura* spp. (Hymenoptera: Xyelidae), on white fir in the central Sierra Nevada of California. III. Mortality factors of egg, larval, and adult stages and a partial life table. Canadian Entomol., 111:833-888.
- Peck, W. B., L. O. Warren, and I. L. Brown. 1971. Spider fauna of shortleaf and loblolly pines in Arkansas. J. Georgia Entomol. Soc., 6:87-94.
- Powell, J. M. 1971. The arthropod fauna collected from the comandra blister rust, *Cronartium comandrae*, on lodgepole pine in Alberta. Canadian Entomol., 103:908-918.
- Price, P. W. 1975. Insect Ecology. John Wiley & Sons, New York, 514 pp.
- Procter, W. 1946. Biological survey of the Mount Desert Region Incorporated. Part VII. Wistar Institute of Anatomy and Biology. Philadelphia., 566 pp.
- Ramsey, H. 1941. Fauna of pine bark. Elisha Mitchell Sci. Soc., 57:91-97.
- Renault, T. R. 1968. An illustrated key to arboreal spiders (Araneida) in the fir-spruce forests of New Brunswick. Canada Dep. Fisheries and Forestry, For. Res. Lab., Fredericton, New Brunswick. Internal Rep. M-39, 41 pp.
- Renault, T. R. and C. A. Miller. 1972. Spiders in a fir-spruce biotype: abundance, diversity, and influence on spruce budworm densities. Canadian J. Zool., 50:1039-1046.

- Stratton, G. E., G. W. Uetz, and D. G. Dillery. 1979. A comparison of the spiders of three coniferous tree species. *J. Arachnol.*, 6:219-226.
- Sokal, R. R. and F. J. Rohlf. 1981. *Biometry*. 2nd ed. W. H. Freeman and Company. New York, 859 pp.
- Taylor, R. L. 1928. The arthropod fauna of coniferous leaders weeviled by *Pissodes strobi* (Peck). *Psyche*, 35:217-225.
- Turnbull, A. L. 1956. Spider predators of the spruce budworm, *Choristoneura fumiferana* (Clem.) at Lillooet, B.C., Canada. *Proc. 8th Pacific Sci. Congr. Vol. IIIA*:1579-1593.
- Worley, L. G. and G. B. Pickwell. 1927. The spiders of Nebraska. *University Studies, University of Nebraska*, 27 (1-4):1-129.

Manuscript received October 1985, revised January 1986.